

CD11c antibody [N418] (Biotin)

Cat. No. GTX74936

Host	Armenian Hamster
Clonality	Monoclonal
Isotype	IgG2
Applications	FCM
Reactivity	Mouse

References (11)

Package

100 µg

PRODUCT

Summary

The N418 antibody reacts with mouse CD11c, also known as integrin α X. This 150 kDa cell surface glycoprotein is part of a family of integrin α receptors that mediate adhesion between cells (cell-cell) and components of the extracellular matrix, e.g. fibrinogen (cell-matrix). In addition, integrin α s are active signaling receptors which recruit leukocytes to inflammatory sites and promote cell activation. Complete, functional integrin α receptors consist of distinct combinations of integrin α chains which are differentially expressed. integrin α X (CD11c) assembles with integrin β 2 (CD18) into a receptor complex known as CR4 which can bind and induce signaling through ICAMs and VCAM-1 on endothelial cells and can also facilitate removal of iC3b bearing foreign cells.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
FCM	Assay dependent

Not tested in other applications.

Properties

Form	Liquid
Buffer	10mM NaH ₂ PO ₄ , 150mM NaCl
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Mouse spleen dendritic cells.
Purification	Purified by affinity chromatography From tissue culture supernatant
Conjugation	Biotin




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

**GTX74936 FCM Image**

FACS analysis of C57Bl/6 splenocytes stained APC MHC Class II antibody, followed by staining with GTX74936 CD11c antibody [N418] (Biotin).

Right panel : Co-stained with APC MHC Class II antibody and GTX74936

Left panel : Co-stained with APC MHC Class II antibody and Biotin Armenian hamster IgG isotype control

Antibody amount : 0.03 μ g



For full product information, images and publications, please visit our [website](#).