

CD68 antibody [514H12]

Cat. No. GTX75390

| | |
|---------------------|-----------------------|
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG2a |
| Applications | ICC/IF, IHC-P, IHC-Fr |
| Reactivity | Human |

Package
50 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| ICC/IF | Assay dependent |
| IHC-P | 1/40-1/80 |
| IHC-Fr | Assay dependent |

Note : This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. EDTA buffer pH 8.0 is recommended for this purpose. N.B. The epitope recognised by clone 514H12 is sensitive to peroxide. It is therefore recommended that

Not tested in other applications.

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Buffer | Tissue culture supernatant |
| Preservative | 0.09% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Immunogen | Fusion protein corresponding to external domain of human CD68. |
| Purification | Tissue Culture Supernatant |
| Conjugation | Unconjugated |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).