

CD200 antibody [MRC OX-2] (PE)

Cat. No. GTX76141

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	FCM, IHC
Reactivity	Rat

Package
50 µg

Applications

Application Note

For FACS: Use at an assay dependent dilution. For IHC: Use at an assay dependent dilution. It is also useful for labeling the follicular dendritic cells thought to be involved in the generation of B cell memory as it does not label the Ia-positive dendritic cells present in the T-dependant areas of lymphoid organs. Optimal dilutions/concentrations should be determined by the researcher.

Product Note

Rat thymocytes, brain, follicular dendritic cells in lymphoid organs, vascular endothelium, and at low levels on some smooth muscle and B lymphocytes

Properties

Form	Liquid
Buffer	PBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
Concentration	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Rat Thymocyte membrane glycoproteins
Purification	Protein G purified
Conjugation	Phycoerythrin (PE) Wavelength


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX76141 FCM Image

Staining of rat thymocytes with Mouse anti Rat CD200 (GTX42296)



For full product information, images and publications, please visit our [website](https://www.genetex.com).