

Macrophage + Monocyte antibody [MAC387]

Cat. No. GTX76577

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-P, IHC-Fr, FCM
Reactivity	Human, Mouse, Rat, Rabbit, Goat, Bovine, Cat, Dog, Guinea pig, Baboon, Cynomolgus monkey, Fallow deer, Horse, Marmoset Pig, Mink, Pygmy hippopotamus, Rhesus Monkey, Squirrel monkey

References (6)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1/100-1/200
IHC-Fr	1/100-1/200
FCM	1/50-1/100

Note : This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase. Permeabilization step is needed.

Not tested in other applications.

Product Note

The antigen recognized by clone MCA387 is expressed by granulocytes, monocytes and by tissue macrophages. Variable results have been reported for staining brain macrophages and microglia.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Human monocytes.
Purification	Protein G purified From tissue culture supernatant
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

Date 2026 / 05 / 23 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).