

## Goat Anti-Chicken IgA antibody (FITC)

## Cat. No. GTX77033

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	IHC-Fr, FCM
Reactivity	Chicken

Package  
500 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	1/200-1/2,000
FCM	Assay dependent

Not tested in other applications.

**Product Note** Goat anti Chicken IgA antibody may react with IgA from other species.

## Properties

Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Purified chicken IgA.
Purification	Purified by affinity chromatography
Conjugation	Fluorescein isothiocyanate (FITC) <a href="#">Wavelength</a>
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 1 of 1