

## BSA antibody (FITC)

## Cat. No. GTX77111

Host	Sheep
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC
Reactivity	Bovine

Package 1 mg

## **Applications**

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2 - 0.05 ug/ml
ICC/IF	10 - 2 ug/ml
IHC	10 -2 ug/ml
Not tested in other applications	

**Calculated MW** 69 kDa. ( <u>Note</u> )

Properties	
Form	Liquid
Buffer	PBS, stabilizer
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Sheep were immunized with highly purified Bovine Serum Albumin in Freund's adjuvant. After multiple immunizations, serum was collected, antibodies were isolated by immunoaffinity chromatography.
Purification	Purified by antigen-affinity chromatography
Conjugation	Fluorescein isothiocyanate (FITC) <u>Wavelength</u> Ratio: 3-6 FITC molecules per IgG molecule.
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.

Date 2025 / 11 / 09 Page 1 of 1