

# Myc tag antibody (FITC)

# Cat. No. GTX77377

Host	Chicken
Clonality	Polyclonal
Isotype	IgY
Applications	WB, ICC/IF, FCM, ELISA, IHC
Reactivity	Species independent

Package 100 μg

# **Applications**

### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2 - 0.05 ug/ml
ICC/IF	10 - 2 ug/ml
FCM	2-10 ug/ml
ELISA	0.1 - 0.01 ug/ml
IHC	Assay dependent

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Chickens were immunized with EQKLISEEDL ( C-MYC) conjugated with KLH. After multiple immunizations in Freunds adjuvant, eggs were collected. Antibodies were immunoaffinity purified using the peptide immobilized on a solid phase. The antibodies were conjugated with Fluorescein Isothiocyanate (FITC).
Purification	Purified by antigen-affinity chromatography
Conjugation	Fluorescein isothiocyanate (FITC) <u>Wavelength</u> Ratio: 3-6 FITC molecules per IgG molecule.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 24 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

### DATA IMAGES



GTX77377 IHC Image



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 24 Page 2 of 2