

Myc tag antibody (FITC)

Cat. No. GTX77382

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC
Reactivity	Species independent

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2 - 0.05 ug/ml
ICC/IF	10 - 2 ug/ml
IHC	10 -2 ug/ml

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Rabbits were immunized with EQKLISEEDL (C-MYC) conjugated with KLH. After multiple immunizations in Freund's adjuvant, serum was collected. Antibodies were immunoaffinity purified using the peptide immobilized on a solid phase
Purification	Purified by antigen-affinity chromatography
Conjugation	Fluorescein isothiocyanate (FITC) Wavelength Ratio : 3-6 FITC molecules per IgG molecule.

Note
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).