

GST tag antibody (FITC)

Cat. No. GTX77388

| | |
|---------------------|---------------------|
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, ICC/IF, IHC |
| Reactivity | Species independent |

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 0.2 - 0.05 ug/ml |
| ICC/IF | 10 - 2 ug/ml |
| IHC | 10 -2 ug/ml |

Not tested in other applications.

Properties

| | |
|----------------------|--|
| Form | Liquid |
| Buffer | PBS, 0.2% BSA |
| Preservative | 0.1% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Glutathione-S-transferase (GST) from Schistosoma japonicum. |
| Purification | Purified by antigen-affinity chromatography |
| Conjugation | Fluorescein isothiocyanate (FITC) Wavelength Ratio : 3-6 FITC molecules per IgG molecule. |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).