

HA tag antibody (FITC)

Cat. No. GTX77397

Host	Chicken
Clonality	Polyclonal
Isotype	IgY
Applications	WB, ICC/IF, ELISA
Reactivity	Species independent

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	10 - 2 ug/ml
ELISA	Assay dependent

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Chickens were immunized with HA-tag conjugated to KLH. After multiple immunizations in Freund's adjuvant, eggs were collected. Antibodies were immunoaffinity purified using the peptide conjugated to a solid-phase support then conjugated with fluorescein i
Purification	Purified by antigen-affinity chromatography
Conjugation	Fluorescein isothiocyanate (FITC) Wavelength Ratio : 3-6 FITC molecules per IgG molecule.
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 1 of 1