

HA tag antibody (FITC)

Cat. No. GTX77397

| | |
|--------------|---------------------|
| Host | Chicken |
| Clonality | Polyclonal |
| Isotype | IgY |
| Applications | WB, ICC/IF, ELISA |
| Reactivity | Species independent |

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | Assay dependent |
| ICC/IF | 10 - 2 ug/ml |
| ELISA | Assay dependent |

Not tested in other applications.

Properties

| | |
|---------------|---|
| Form | Liquid |
| Buffer | PBS, 0.2% BSA |
| Preservative | 0.1% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Chickens were immunized with HA-tag conjugated to KLH. After multiple immunizations in Freund's adjuvant, eggs were collected. Antibodies were immunoaffinity purified using the peptide conjugated to a solid-phase support then conjugated with fluorescein i |
| Purification | Purified by antigen-affinity chromatography |
| Conjugation | Fluorescein isothiocyanate (FITC) Wavelength Ratio : 3-6 FITC molecules per IgG molecule. |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).