

Claudin 17 antibody, Internal

Cat. No. GTX77836

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 ug/ml
IHC-P	2-10 ug/ml

Not tested in other applications.

Calculated MW 25 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to an Internal region of Human Claudin 17
Purification	Protein A purified
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX77836 IHC-P Image

IHC-P analysis of human spleen tissue using GTX77836 Claudin 17 antibody at 4.0-8.0µg/ml.


GTX77836 IHC-P Image

IHC-P analysis of human lung tissue using GTX77836 Claudin 17 antibody at 4.0-8.0µg/ml.


GTX77836 WB Image

WB analysis of Jurkat cells using GTX77836 Claudin 17 antibody at 2.0µg/ml.



For full product information, images and publications, please visit our [website](https://www.genetex.com).