

## NTAL antibody [NAP-07] (PE)

**Cat. No. GTX78366**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	ICC/IF, FCM
<b>Reactivity</b>	Human, Mouse

**Package**  
100 µg

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
FCM	Assay dependent

Not tested in other applications.

**Product Note** This antibody reacts with Non-T cell activation linker (NTAL).

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS
<b>Preservative</b>	15mM Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
<b>Concentration</b>	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Recombinant cytoplasmic domain (aa 91-243) of human NTAL.
<b>Purification</b>	Purified by size-exclusion chromatography
<b>Conjugation</b>	Phycoerythrin (PE) <a href="#">Wavelength</a>


**Note** For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

## DATA IMAGES

**GTX78366 ICC/IF Image**

Subcellular localization of NTAL by confocal microscopy in THP-1 human acute monocytic leukemia cell line. Subcellular localization of NTAL by confocal microscopy in THP-1 human acute monocytic leukemia cell line. THP-1 cells were permeabilized and immunostained using anti-NTAL (NAP-07 (GTX23992); green). Nuclei are visualized by propidium iodide (red).



For full product information, images and publications, please visit our [website](#).