

NTAL antibody [NAP-07] (PE)

Cat. No. GTX78366

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	ICC/IF, FCM
Reactivity	Human, Mouse

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
FCM	Assay dependent

Not tested in other applications.

Product Note This antibody reacts with Non-T cell activation linker (NTAL).

Properties

Form	Liquid
Buffer	PBS
Preservative	15mM Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
Concentration	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant cytoplasmic domain (aa 91-243) of human NTAL.
Purification	Purified by size-exclusion chromatography
Conjugation	Phycoerythrin (PE) Wavelength


Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX78366 ICC/IF Image

Subcellular localization of NTAL by confocal microscopy in THP-1 human acute monocytic leukemia cell line. Subcellular localization of NTAL by confocal microscopy in THP-1 human acute monocytic leukemia cell line. THP-1 cells were permeabilized and immunostained using anti-NTAL (NAP-07 (GTX23992); green). Nuclei are visualized by propidium iodide (red).



For full product information, images and publications, please visit our [website](#).