CD10 antibody [LT10] (FITC)

Cat. No. GTX78376

| Host | Mouse |
|-------------|------------|
| Clonality | Monoclonal |
| lsotype | lgG1 |
| Application | FACS |
| Reactivity | Human |

Package

100 test

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|-----------------------------------|---|
| FACS | 20 μl reagent / 100 μl of whole blood or 10^6 cells in a suspension |
| Not tested in other applications. | |

Calculated MW

86 kDa. (<u>Note</u>)


| PROPERTIES | |
|---------------|---|
| Form | Liquid |
| Buffer | PBS |
| Preservative | 15mM Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light. |
| Concentration | Batch dependent (Please refer to the vial label for the specific concentration.) |
| Immunogen | mouse NALM-6 leukemia pre-B cell line (tissue/cell preparation) |
| Purification | Purified by size-exclusion chromatography |
| Conjugation | Fluorescein isothiocyanate (FITC) |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our <u>website</u>.


GeneTex International Corporation (Global) € 886-3-6208988
В 886-3-6208989
S infoasia@genetex.com

DATA IMAGES



GTX78376 FACS Image

FACS analysis of human CD10 positive Neutrophil Granulocytes (red-filled) and human CD10 negative Lymphocytes (black-dashed) using GTX78376 CD10 antibody [LT10] (FITC). Antibody amount : 20μ l reagent / 100μ l of peripheral whole blood



GTX78376 FACS Image

FACS analysis of human peripheral blood using GTX78376 CD10 antibody [LT10] (FITC). Antibody amount : 20 μ l reagent / 100 μ l of peripheral whole blood



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 14 Page 2 of 2