

# Cytochrome P450 2C11 antibody

**Cat. No. GTX79174**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Application</b>	WB, ICC/IF, IHC-P, IHC-Fr, ELISA
<b>Reactivity</b>	Human, Rat, Cat

**Package**  
100 µl

## APPLICATION

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1,500
ICC/IF	1:20-1:200
IHC-P	1:100-1:500
IHC-Fr	1:100
ELISA	Assay dependent

Not tested in other applications.

**Calculated MW** 57 kDa. ( [Note](#) )

## PROPERTIES

<b>Form</b>	Liquid
<b>Buffer</b>	Serum diluted with PBS
<b>Preservative</b>	0.05% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Immunogen</b>	Synthetic peptides corresponding to residues D(49) I G Q S I K K F S K V(60) and Q(491) R A D S L S H L(500) of rat Cytochrome P450 2C11.
<b>Purification</b>	Unpurified
<b>Conjugation</b>	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

**DATA IMAGES**

**GTX79174 ICC/IF Image**

ICC/IF analysis of PC-12 cells using GTX79174 Cytochrome P450 2C11 antibody. Cells were probed without (left) or with(right) an antibody.


Green : Primary antibody

Blue : Nuclei

Fixation : Formalin

Permeabilization : 0.1% Triton X-100 in TBS for 5-10 minute

Dilution : 1:100 incubated overnight at 4 °C


**GTX79174 ICC/IF Image**

ICC/IF analysis of HeLa cells using GTX79174 Cytochrome P450 2C11 antibody. Cells were probed without (left) or with(right) an antibody.

Green : Primary antibody


Blue : Nuclei

Red : Actin

Fixation : Formalin

Permeabilization : 0.1% Triton X-100 in TBS for 5-10 minute

Dilution : 1:100 incubated overnight at 4 °C


**GTX79174 ICC/IF Image**

ICC/IF analysis of H-4-II-E cells using GTX79174 Cytochrome P450 2C11 antibody. Cells were probed without (left) or with(right) an antibody.

Green : Primary antibody

Blue : Nuclei

Red : Actin

Fixation : Formalin

Permeabilization : 0.1% Triton X-100 in TBS for 5-10 minute

Dilution : 1:100 incubated overnight at 4 °C



For full product information, images and publications, please visit our [website](#).