

TCR V beta 8a antibody [I6G8]

Cat. No. GTX79236

Host	Mouse	
Clonality	Monoclonal	
Isotype	lgG2b	
Applications	WB, ICC/IF, IHC-Fr, FCM, IP, Activation, IHC	
Reactivity	Human, Primate	

Package 200 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

WB Assay dependent ICC/IF 2 μg/ml IHC-Fr Assay dependent FCM Assay dependent IP Assay dependent Activation Assay dependent	Suggested dilution	Recommended dilution
IHC-Fr Assay dependent FCM Assay dependent IP Assay dependent Activation Assay dependent	WB	Assay dependent
FCM Assay dependent IP Assay dependent Activation Assay dependent	ICC/IF	2 μg/ml
IP Assay dependent Activation Assay dependent	IHC-Fr	Assay dependent
Activation Assay dependent	FCM	Assay dependent
	IP	Assay dependent
IHC Assay dependent	Activation	Assay dependent
Assay dependent	IHC	Assay dependent

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS, 0.5% BSA, Glycerol
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.15 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Human TCR V beta 8
Purification	Protein A purified
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 12 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX79236 ICC/IF Image

ICC/IF analysis of MOLT-4 cells using GTX79236 TCR V beta 8a antibody [I6G8]. Panel e is a no primary antibody control.

Green: Primary antibody

Blue: Nuclei Red: Actin

Fixation: 4% paraformaldehyde

Permeabilization: 0.1% Triton X-100 for 10 minute

Dilution: 2 µg/ml in 0.1% BSA incubated for 3 hours at room temperature



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 12 Page 2 of 2