

## CD86 antibody [BU63] (APC)

## Cat. No. GTX80210

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	FCM
Reactivity	Human, Dog

References ( 5 )

Package

100 test

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
FCM	10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension
Not tested in other applications.	

**Product Note** Domain depletion epitope mapping studies indicate that the binding site of Mouse anti Human CD86, clone Bu63 is located within the Ig-v-like domain of human CD86 (Jeanin et al. 1997).

## Properties


Form	Liquid
Buffer	PBS
Preservative	15mM Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	B-lymphoblastoid cell line ARH 77
Purification	Purified by size-exclusion chromatography
Conjugation	Allophycocyanin (APC) <a href="#">Wavelength</a>
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 16 Page 1 of 2

## DATA IMAGES



## GTX80210 FCM Image

FACS analysis of human peripheral blood using GTX80210 CD86 antibody [BU63] (APC).



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 16 Page 2 of 2