

# HLA-G antibody [87G] (Low Endotoxin)

# Cat. No. GTX80252

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2a
Applications	IHC-Fr, FCM, ELISA, Functional Assay
Reactivity	Human

References ( 1 ) Package 100 μg

# Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-Fr	Assay dependent
FCM	Assay dependent
ELISA	Assay dependent
Functional Assay	Assay dependent

Not tested in other applications.

**Product Note** 

This antibody recognizes both membrane-bound and soluble forms of HLA-G (HLA-G1 and HLA-G5). The antibody 87G blocks interaction of HLA-G with inhibitory receptors.

We do not recommend use of this product for Mouse, Rat samples.

Form       Liquid         Buffer       Filter-sterilized PBS         Preservative       No preservatives         Storage       Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.         Concentration       1 mg/ml (Please refer to the vial label for the specific concentration.)         Immunogen       HLA-B27 transgenic mice were imunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.         Purification       Protein A purified         Endotoxin       < 0.01 EU/μg (determined by the LAL assay)         Conjugation       Unconjugated	Properties	
Preservative       No preservatives         Storage       Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.         Concentration       1 mg/ml (Please refer to the vial label for the specific concentration.)         Immunogen       HLA-B27 transgenic mice were imunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.         Purification       Protein A purified         Endotoxin       < 0.01 EU/μg (determined by the LAL assay)	Form	Liquid
Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.  Concentration 1 mg/ml (Please refer to the vial label for the specific concentration.)  Immunogen HLA-B27 transgenic mice were imunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.  Purification Protein A purified  Endotoxin < 0.01 EU/µg (determined by the LAL assay)	Buffer	Filter-sterilized PBS
Concentration       1 mg/ml (Please refer to the vial label for the specific concentration.)         Immunogen       HLA-B27 transgenic mice were imunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.         Purification       Protein A purified         Endotoxin       < 0.01 EU/μg (determined by the LAL assay)	Preservative	No preservatives
Immunogen       HLA-B27 transgenic mice were imunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.         Purification       Protein A purified         Endotoxin       < 0.01 EU/μg (determined by the LAL assay)	Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
HLA-G and human beta2-microglobulin.  Purification Protein A purified  Endotoxin < 0.01 EU/μg (determined by the LAL assay)	Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Endotoxin < 0.01 EU/μg (determined by the LAL assay)	Immunogen	
	Purification	Protein A purified
Conjugation Unconjugated	Endotoxin	< 0.01 EU/μg (determined by the LAL assay)
	Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 12 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

### DATA IMAGES



### **GTX80252 Functional Assay Image**

Effect on cytolytical activity of human polyclonal NK cells on M8 cells transfected with empty vector (column 1) and with HLA-G1 cDNA (columns 2-6). Blocking of HLA-G1 on transfectants with anti-human HLA-G (87G) restored specific lysis.

Blocking antibodies -

Column 1-2: None

Column 3: Isotype mouse IgG2a control

Column 4: GTX80252 anti-human HLA-G (87G) purified Column 5: anti-human HLA-G (87G) F(ab')2 fragment

Column 6: anti-human HLA-G (MEM-G/9)



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 12 Page 2 of 2