

DEK antibody, C-term

Cat. No. GTX80509

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, FCM
Reactivity	Human

Package
400 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000
ICC/IF	1:10-1:50
IHC-P	1:50-1:100
FCM	1:10-1:50

Not tested in other applications.

Calculated MW 43 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	KLH conjugated synthetic peptide between 336-364 amino acids from the C-terminal region of human DEK.
Purification	Protein A purified, followed by peptide affinity purification.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTx80509 WB Image

WB analysis of U-251 MG and PC-3 cell using GTx80509 DEK antibody, C-term.

Loading : 35ug per lane


Dilution : 1:1000


GTx80509 FCM Image

FACS analysis of WiDr cells using GTx80509 DEK antibody, C-term.

Green : primary antibody

Blue : negative control


GTx80509 IHC-P Image

IHC-P analysis of human testis tissue using GTx80509 DEK antibody, C-term.



For full product information, images and publications, please visit our [website](http://www.genetex.com).