

LAG3 antibody, Internal

Cat. No. GTX81605

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, FCM
Reactivity	Human, Mouse

Package
400 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000
IHC-P	1:10-1:50
FCM	1:25

Note : Permeabilization : 90% methanol for 10 min

Not tested in other applications.

Calculated MW 57 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	KLH conjugated synthetic peptide between 103-132 amino acids from the Central region of human LAG3.
Purification	Protein A purified, followed by peptide affinity purification.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

**GTX81605 WB Image**

WB analysis of various samples using GTX81605 LAG3 antibody, Internal.


Lane 1: K562 whole cell lysates

Lane 2: human liver lysates

Lane 3: human ovary lysates

Loading : 20 µg per lane

Dilution : 1:2000

**GTX81605 WB Image**


WB analysis of various samples using GTX81605 LAG3 antibody, Internal.

Lane 1: human ovary lysate

Lane 2: mouse liver lysate

Loading : 20 µg per lane

Dilution : 1:1000

**GTX81605 WB Image**

WB analysis of mouse heart lysate using GTX81605 LAG3 antibody, Internal.

Loading : 20 µg per lane

Dilution : 1:2000



For full product information, images and publications, please visit our [website](#).