

ABCG1 antibody, Internal

Cat. No. GTX81867

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, FCM
Reactivity	Human

References (1)

Package

400 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000
IHC-P	1:10-1:50
FCM	1:10-1:50

Not tested in other applications.

Calculated MW 76 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	KLH conjugated synthetic peptide between 359-387 amino acids from the Central region of human ABCG1.
Purification	Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Conjugation	Unconjugated

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX81867 WB Image


WB analysis of HepG2 whole cell lysate using GTX81867 ABCG1 antibody, Internal.

Loading : 20 µg per lane

Dilution : 1:1000


GTX81867 IHC-P Image

IHC-P analysis of human hepatocarcinoma using GTX81867 ABCG1 antibody, Internal.


GTX81867 FCM Image

FACS analysis of HepG2 cells using GTX81867 ABCG1 antibody, Internal.

Top histogram : negative control

Bottom histogram : HepG2 cells



For full product information, images and publications, please visit our [website](https://www.genetex.com).