

Coxsackie Adenovirus Receptor antibody, Internal

Cat. No. GTX81978

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, FCM
Reactivity	Human

Package
400 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000
FCM	1:10-1:50

Not tested in other applications.

Calculated MW 40 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	KLH conjugated synthetic peptide between 132-161 amino acids from the Central region of human CXADR.
Purification	Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX81978 FCM Image

FACS analysis of NCI-H460 cells using GTX81978 Coxsackie Adenovirus Receptor antibody, Internal.

Top histogram : negative control

Bottom histogram : NCI-H460 cells


GTX81978 WB Image

WB analysis of NCI-H460 cell lysate (35ug/lane) using GTX81978 Coxsackie Adenovirus Receptor antibody, Internal.



For full product information, images and publications, please visit our [website](https://www.genetex.com).