

EIF4E2 antibody

Cat. No. GTX82524

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse

References (7)

Package

400 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:2000
ICC/IF	Assay dependent
IHC-P	1:10-1:50

Not tested in other applications.

Calculated MW 28 kDa. ([Note](#))


Properties

Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	This EIF4E2 antibody is generated from rabbits immunized with human EIF4E2 recombinant protein.
Purification	Protein A purified, followed by peptide affinity purification.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


DATA IMAGES



GTX82524 WB Image


WB analysis of various samples using GTX82524 EIF4E2 antibody.

- Lane 1: 293 whole cell lysate
- Lane 2: HeLa whole cell lysate
- Lane 3: HL-60 whole cell lysate
- Lane 4: RD whole cell lysate
- Lane 5: C2C12 whole cell lysate
- Loading : 20 µg per lane
- Dilution : 1:2000



GTX82524 IHC-P Image

IHC-P analysis of human breast carcinoma using GTX82524 EIF4E2 antibody.



GTX82524 WB Image

WB analysis of various samples using GTX82524 EIF4E2 antibody.

- Lane 1: 293 whole cell lysate
- Lane 2: RD whole cell lysate
- Lane 3: HeLa whole cell lysate
- Lane 4: HepG2 whole cell lysate
- Lane 5: HL-60 whole cell lysate
- Loading : 20 µg per lane
- Dilution : 1:2000



For full product information, images and publications, please visit our [website](#).