

# Peroxidase Enhancer

**Cat. No. GTX82979**

**Applications** ICC/IF, IHC

References ( 2 )  
 Package  
 1 kit

## PRODUCT

**Summary** Peroxidase Signal Enhancer

## Applications

### Application Note

Mixing of reagent: To five ml buffer (reagent B) in a test tube, add one drop of concentrated Enhancing reagent (reagent E) mix well. This ready to use reagent is good for 4-6 hours. Procedure: 1.) After the peroxidase reaction, wash slide with buffer (this buffer should not contain any sodium azide because peroxidase is inactivated) and distilled/ deionized water. 2.) Add few drops of ready to use enhancing reagent for 1-2 minutes. 3.) Add few drops of ready to use AEC reagent, or DAB reagent, follow instructions as outlined for these chromogens.

## Properties

**Form** Liquid

**Buffer** Reagent B (buffer): 250ml Reagent E (conc. Enhancing reagent): 3 ml

**Preservative** No preservatives

**Storage** Keep as concentrated solution, aliquot and store at 4°C. DO NOT FREEZE.

**Note** For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.  
 Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).