

CBWD1 antibody [4E8]

Cat. No. GTX84728

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2b
Applications	WB, IHC-P, FCM
Reactivity	Human, Rat, Dog, Monkey

Package 100 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500
IHC-P	Assay dependent
FCM	1:100
Not tested in other applications	

Not tested in other applications.

Calculated MW 44 kDa. (Note)


Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.6 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length human recombinant protein of human CBWD1 (NP_060961) produced in HEK293T cell.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 14 Page 1 of 2

DATA IMAGES




GTX84728 FCM Image

FACS analysis of Jurkat cells using GTX84728 CBWD1 antibody [4E8].

Red: Primary antibody


Blue: Negative control antibody



GTX84728 WB Image


WB analysis of various cell lines using GTX84728 CBWD1 antibody [4E8].

Loading: 35 ug per lane



GTX84728 IHC-P Image

IHC-P analysis of human lymph node tissue using GTX84728 CBWD1 antibody [4E8].



GTX84728 WB Image

WB analysis of HEK293T cells transfected with CBWD1 plasmid (Right) or empty vector (Left) for 48 hrs using GTX84728 CBWD1 antibody [4E8].

Loading: 5 ug per lane



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 14 Page 2 of 2