

ADSL antibody [2D10]

Cat. No. GTX84956

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, IHC-P, FCM
Reactivity	Human, Rat, Dog, Monkey

References (2)

Package

100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:2000
IHC-P	1:150
FCM	1:100

Not tested in other applications.

Calculated MW 55 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS, 1% BSA, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.55 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length human recombinant protein of human ADSL (NP_000017) produced in HEK293T cell.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 1 of 2

DATA IMAGES


**GTX84956 FCM Image**

FACS analysis of HEK293T cells transfected with either ADSL plasmid(Red) or empty vector control plasmid(Blue) using GTX84956 ADSL antibody [2D10].

**GTX84956 WB Image**


WB analysis of HEK293T cells transfected with ADSL plasmid (Right) or empty vector (Left) for 48 hrs using GTX84956 ADSL antibody [2D10].

Loading : 5 ug per lane

**GTX84956 IHC-P Image**

IHC-P analysis of human prostate carcinoma tissue using GTX84956 ADSL antibody [2D10].

Antigen retrieval : Heat-induced epitope retrieval by 10mM citrate buffer, pH6.0, 100°C for 10min.

**GTX84956 IHC-P Image**

IHC-P analysis of human kidney carcinoma tissue using GTX84956 ADSL antibody [2D10].

Antigen retrieval : Heat-induced epitope retrieval by 10mM citrate buffer, pH6.0, 100°C for 10min.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 2 of 2