

MD1 antibody

Cat. No. GTX85066

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.5 - 2 µg/mL
IHC-P	2 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 18 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	MD-1 antibody was raised against a 14 amino acid synthetic peptide from near the carboxy terminus of human MD-1. The immunogen is located within the last 50 amino acids of MD-1.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX85066 IHC-P Image

IHC-P analysis of human spleen tissue using GTX85066 MD1 antibody.
Working concentration : 2 µg/ml



GTX85066 WB Image

WB analysis of Daudi cell lysate using GTX85066 MD1 antibody.
Working concentration : (A) 0.5, (B) 1, and (C) 2 µg/ml



For full product information, images and publications, please visit our [website](#).