

TIM-1 antibody

Cat. No. GTX85068

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Human, Mouse

References (2)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 µg/mL
IHC-P	10 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 39 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	TIM-1 antibody was raised against a 16 amino acid synthetic peptide from near the amino terminus of human TIM-1. The immunogen is located within amino acids 50 - 100 of TIM-1.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 08 Page 1 of 2


DATA IMAGES



GTX85068 WB Image

WB analysis of 3T3 cell lysate using GTX85068 TIM-1 antibody.

Working concentration : 1 μ g/ml




GTX85068 WB Image

WB analysis of NIH/3T3 cell lysates using GTX85068 TIM-1 antibody.

Loading : 15 μ g of lysates per lane


Dilution : 8 μ g/ml



GTX85068 IHC-P Image

IHC-P analysis of human uterus tissue using GTX85068 TIM-1 antibody.

Working concentration : 10 μ g/ml



GTX85068 WB Image

WB analysis of various sample lysates using GTX85068 TIM-1 antibody.

Loading : 15 μ g of lysates per lane

Dilution : 8 μ g/ml



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 08 Page 2 of 2