

## RASD2 antibody

Cat. No. GTX85427

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Human, Mouse, Rat

Package  
100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 - 2 µg/mL
IHC-P	2.5 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW	30 kDa. ( <a href="#">Note</a> )
Product Note	TEM2 antibody may cross-react with the closely related protein RASD1.

## Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	RASD2 antibody was raised against a 16 amino acid synthetic peptide near the amino terminus of the human RASD2. The immunogen is located within amino acids 20 - 70 of RASD2.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

## DATA IMAGES




### GTx85427 WB Image

WB analysis of rat colon tissue lysate using GTx85427 RASD2 antibody.  
Working concentration : (A) 1 and (B) 2 µg/ml



### GTx85427 IHC-P Image

IHC-P analysis of human colon tissue using GTx85427 RASD2 antibody.  
Working concentration : 20 µg/ml



### GTx85427 IHC-P Image

IHC-P analysis of human colon tissue using GTx85427 RASD2 antibody.  
Working concentration : 2.5 µg/ml



For full product information, images and publications, please visit our [website](#).