

FNTA antibody

Cat. No. GTX85881

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse, Rat, Dog, Chicken

References (1) Package 50 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 - 3 μg/ml
IHC-P	10 μg/ml
No constitution of the con	

Not tested in other applications.

Calculated MW 44 kDa. (Note)


Properties	
Form	Liquid
Buffer	PBS, 0.2% Gelatin
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A portion of amino acids 150-250 of human farnesyltransferase.
Purification	Protein G purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 13 Page 1 of 2

DATA IMAGES




GTX85881 IHC-P Image

IHC-P analysis of human prostate tissue using GTX85881 FNTA antibody.

Antigen retrieval: Heat-induced antigen retrieval

Dilution: 10 μg/ml



GTX85881 WB Image

WB analysis of mouse colon cell lysate in the 1) absence and 2) presence of immunizing peptide using GTX85881 FNTA antibody.

Dilution : $2 \mu g/ml$



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 13 Page 2 of 2