

## Streptavidin (FITC)

## Cat. No. GTx85911

## Applications

ICC/IF, IHC-P, FCM

## Package

500 µg

## PRODUCT

## Summary

Streptavidin is a 55 kDa (subunit MW 14 kDa) biotin-binding protein isolated from *Streptomyces avidini*. Streptavidin is superior to avidin, because it does not contain carbohydrate like avidin and has no net charge at neutral pH.

Streptavidin~biotin system is routinely used in Immunohistochemistry (IHC). Extinction Coefficient 1% A280=32.0;

Eletrophoretically homogenous, single band. Protein content: > 95%; One mg of Streptavidin is conjugated with FITC, unconjugated Streptavidin is removed. This conjugate of streptavidin is recommended for use with Biotin~conjugated antibodies. Avidin~Biotin conjugated reagents are stable, sensitive and give less background. Fluorophore: Fluorescein isothiocyanate A max=492nm; E=520nm; Fluorophore/Protein: A492nm/A280nm=~1

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-P	Assay dependent
FCM	Assay dependent

Not tested in other applications.

## Properties

## Note

Form	Liquid
Buffer	10mM Phosphate, 150mM NaCl, 1% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
Conjugation	Fluorescein isothiocyanate (FITC) <a href="#">Wavelength</a>
	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 28 Page 1 of 1