

Streptavidin (HRP)

Cat. No. GTX85912

Applications

WB, ELISA, IHC

References (6)

Package

500 µg

PRODUCT

Summary

Streptavidin is a 22- 55 kDa (subunit MW 14 kDa) biotin-binding protein isolated from Streptomyces avidini. Streptavidin is superior to avidin, because it does not contain carbohydrate like avidin and has no net charge at neutral pH. Streptavidin~biotin system is routinely used in Immunohistochemistry (IHC). Extinction Coefficient 1% A280=32.0. Protein content: > 95%; Peroxidase used for conjugation is 44 kDa glycoprotein isolated from horseradish roots. The Rz (Reinheitszahl), absorbance ratio of A403/A275 is ~3.0. One mg of Streptavidin is conjugated with peroxidase, in equimolar ratio. Unconjugated Streptavidin is removed.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution

Recommended dilution

WB Assay dependent

ELISA Assay dependent

IHC Assay dependent

Not tested in other applications.

Properties

Form Liquid

Buffer 10mM Phosphate, 200mM NaCl, 1% BSA

Preservative 0.05% ProClin 300

Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.

Concentration 1 mg/ml (Please refer to the vial label for the specific concentration.)

Conjugation Horseradish peroxidase(HRP)

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).