

OATP8 antibody, Internal

Cat. No. GTX88295

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, FCM
Reactivity	Human

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
FCM	Assay dependent

Not tested in other applications.

Calculated MW 77 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-NGTSPEIVEKD, from the internal region of the protein sequence according to NP_062818.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 1 of 2

DATA IMAGES



GTX88295 FCM Image


FACS analysis of PFA-fixed HepG2 cells using GTX88295 OATP8 antibody, Internal.

Blue : Primary antibody

Black : IgG control

Permeabilization : 0.5% Triton

Dilution : 10 μ g/ml



GTX88295 ICC/IF Image


ICC/IF analysis of PFA-fixed A431 cells using GTX88295 OATP8 antibody, Internal.

Green : Primary antibody

Blue : DAP Negative control: Unimmunized goat IgG (10 μ g/ml)

Permeabilization : 0.15% Triton

Dilution : 10 μ g/ml



GTX88295 WB Image

WB analysis of HEK293 (A) and A431 (B) cell lysate (35 μ g protein in RIPA buffer) using GTX88295 OATP8 antibody, Internal.

Dilution : 1 μ g/ml



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 2 of 2