

Tankyrase 2 antibody, Internal

Cat. No. GTX88516

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.5-1.5µg/ml

Not tested in other applications.

Calculated MW	127 kDa. (Note)
Product Note	Immunogen shows no cross reactivity to TANK1.

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Fusion protein containing 73 aas from aa800-900 region of the protein sequence according to NP_079511.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX88516 WB Image

WB analysis of HEK293 overexpressing Tankyrase2 (lane 4) and TANK1 (lane 2) (mock transfection in first lane) using GTX88516 Tankyrase 2 antibody, Internal. Lower panel shows the same lysates probed for alpha-Actin to show protein levels.

Dilution : 0.5µg/ml



For full product information, images and publications, please visit our [website](https://www.genetex.com).