

Monoglyceride lipase antibody, Internal (near N-term)

Cat. No. GTX88653

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.5-1.5µg/ml
IHC-P	5µg/ml

Note : Human Liver, expression of MGLL is detected in hepatocytes.

Not tested in other applications.

Calculated MW	33 kDa. (Note)
Product Note	This antibody is expected to recognize both reported isoforms (NP_009214.1; NP_001003794.1).

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-QDPLHLVNADGQY, from the internal region (near N Terminus) of the protein sequence according to NP_009214.1; NP_001003794.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).


Date 2026 / 10 / 13 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX88653 IHC-P Image

IHC-P analysis of human liver using GTX88653 Monoglyceride lipase antibody, Internal (near N-term).

Antigen retrieval : citrate buffer pH 6

Dilution : 5µg/ml



GTX88653 WB Image

WB analysis of human frontal cortex (A) and mouse adipose (B) lysates using GTX88653 Monoglyceride lipase antibody, Internal (near N-term).

Dilution : 0.5µg/ml

Loading : 35µg protein in RIPA buffer



For full product information, images and publications, please visit our [website](https://www.genetex.com).