

Neuroligin 3 antibody, Internal

Cat. No. GTX88872

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.05-0.2µg/ml

Not tested in other applications.

Calculated MW 94 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-RKDKRRQEPLRQPS, from the internal region of the protein sequence according to NP_061850.2.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX88872 WB Image

WB analysis of human brain (amygdala) lysate using GTX88872 Neuroligin 3 antibody, Internal.

Dilution : 0.1µg/ml

Loading : 35µg protein in RIPA buffer


GTX88872 WB Image

WB analysis of HEK293 overexpressing Neuregulin3 (mock transfection in first lane) using GTX88872 Neuroligin 3 antibody, Internal.



For full product information, images and publications, please visit our [website](https://www.genetex.com).