

## ARSB antibody, Internal

## Cat. No. GTX88904

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

## Package

100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.03-0.1µg/ml
IHC-P	5-10µg/ml

**Note : Human Skeletal Muscle shows striated pattern staining of majority of muscular fibres.**

Not tested in other applications.

**Calculated MW** 60 kDa. ([Note](#))

**Product Note** This antibody is expected to recognise both reported isoforms (NP\_000037.2; NP\_942002.1).

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	TBS, 0.5% BSA
<b>Preservative</b>	0.02% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Peptide with sequence C-KLARGHTNGTKPLD, from the internal region of the protein sequence according to NP_000037.2; NP_942002.1.
<b>Purification</b>	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
<b>Conjugation</b>	Unconjugated



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 13 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


## DATA IMAGES

**GTX88904 IHC-P Image**

IHC-P analysis of human skeletal muscle using GTX88904 ARSB antibody, Internal.

Antigen retrieval : citrate buffer pH 6

Dilution : 5 $\mu$ g/ml

**GTX88904 WB Image**

WB analysis of human heart lysate using GTX88904 ARSB antibody, Internal.

Dilution : 0.03 $\mu$ g/ml

Loading : 35 $\mu$ g protein in RIPA buffer



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 2 of 2