

SMEK1 antibody, Internal

Cat. No. GTX88983

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.3-1µg/ml

Not tested in other applications.

Calculated MW 95 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-EQQRERQDNPKLD, from the internal region of the protein sequence according to NP_115949.3.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX88983 WB Image

WB analysis of human pancreas lysate using GTX88983 SMEK1 antibody, Internal.

Dilution : 0.3µg/ml

Loading : 35µg protein in RIPA buffer


GTX88983 WB Image

WB analysis of HEK293 overexpressing SMEK1 (mock transfection in first lane) using GTX88983 SMEK1 antibody, Internal.



For full product information, images and publications, please visit our [website](https://www.genetex.com).