

SNAP25 antibody, C-term

Cat. No. GTX89577

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, IHC
Reactivity	Human, Mouse

References (4)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.01-0.03µg/ml
IHC-P	1-2µg/ml
IHC	Assay dependent

Note : Human Cerebellum shows strong axon staining, particularly in the granular region.

Not tested in other applications.

Calculated MW	23 kDa. (Note)
Product Note	This antibody is expected to recognise both reported isoforms NP_003072.2 (SNAP25A) and NP_570824.1 (SNAP25B).

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-DEANQRATKMLGSG, from the C Terminus of the protein sequence according to NP_003072.2; NP_570824.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX89577 IHC-P Image

IHC-P analysis of human cerebellum using GTX89577 SNAP25 antibody, C-term.

Antigen retrieval : citrate buffer Ph 6

Dilution : 1µg/ml



GTX89577 WB Image

WB analysis of human brain (cerebellum) (A) and mouse brain (B) lysates using GTX89577 SNAP25 antibody, C-term.

Dilution : 0.03µg/ml

Loading : 35µg protein in RIPA buffer



For full product information, images and publications, please visit our [website](#).