

CREM antibody, C-term

Cat. No. GTX89673

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.3-1µg/ml
Not tested in other applications.	

Calculated MW 37 kDa. ([Note](#))

Product Note

This antibody is expected to recognise the following protein isoforms of human CREM protein: a (NP_853549.1); e (NP_874387.1); f (NP_874388.1); k (NP_874393.1); l (NP_874394.1); m (NP_877570.1); n (NP_877571.1); o (NP_877572.1); p (NP_877573.1); q (NP_878270.1); v (NP_898883.1). This antibody is not expected to cross-react with CREB protein.

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence LETLKDICSPKTDY, from the C Terminus of the protein sequence according to NP_853549.1; NP_874387.1; NP_874388.1; NP_874393.1; NP_874394.1; NP_877570.1; NP_877571.1; NP_877572.1; NP_877573.1; NP_878270.1; NP_898883.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).


Date 2026 / 10 / 06 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX89673 WB Image

WB analysis of human heart lysate using GTX89673 CREM antibody, C-term.

Dilution : 1µg/ml

Loading : 35µg protein in RIPA buffer



For full product information, images and publications, please visit our [website](#).