

KLF15 antibody, N-term

Cat. No. GTX89919

Host	Goat	Package
Clonality	Polyclonal	100 µg
Isotype	IgG	
Applications	WB, FCM	
Reactivity	Human	

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.3-1µg/ml
FCM	10µg/ml

Not tested in other applications.

Calculated MW 44 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence VDHLLPVDENFSS-C, from the N Terminus of the protein sequence according to NP_054798.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 02 Page 1 of 2

DATA IMAGES




GTX89919 WB Image

WB analysis of HepG2 nuclear (A) and HeLa (B) cell lysate using GTX89919 KLF15 antibody, N-term.

Dilution : 1 μ g/ml

Loading : 35 μ g protein in RIPA buffer



GTX89919 FCM Image

FACS analysis of PFA fixed HepG2 cells using GTX89919 KLF15 antibody, N-term.

Blue : Primary antibody

Black : Isotype control

Permeabilization : 0.5% Triton

Dilution : 10 μ g/ml



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 02 Page 2 of 2