

DKK3 antibody, C-term

Cat. No. GTX89935

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.03-0.1 µg/ml
IHC-P	5-10 µg/ml

Note : Human Brain Cortex shows vesiculate staining in cytoplasm of neuronal cell bodies.

Not tested in other applications.

Calculated MW 38 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-AAAALLGGEI, from the C Terminus of the protein sequence according to NP_037385.2; NP_056965.3; NP_001018067.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX89935 IHC-P Image

IHC-P analysis of human cortex using GTX89935 DKK3 antibody, C-term.

Antigen retrieval : citrate buffer pH 6

Dilution : 5µg/ml



GTX89935 WB Image

WB analysis of HEK293 lysate using GTX89935 DKK3 antibody, C-term.

Dilution : 1µg/ml

Loading : 30µg protein in RIPA buffer



For full product information, images and publications, please visit our [website](https://www.genetex.com).