

## ELMO1 antibody, C-term

## Cat. No. GTX89955

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IP
Reactivity	Human, Mouse, Rat, Zebrafish

## Package

100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.3µg/ml
IP	Assay dependent

Not tested in other applications.

Calculated MW	84 kDa. ( <a href="#">Note</a> )
Product Note	This antibody is expected to recognise both human isoforms of this protein. NP_569709.1 and NP_001034548.1 represent the same isoform.

## Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence PKEPSNYDFVYDCN, from the C Terminus of the protein sequence according to NP_055615.8; NP_569709.1; NP_001034548.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 03 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

**DATA IMAGES****GTX89955 WB Image**

WB analysis of human frontal cortex (A), mouse (B) and rat (C) brain lysate using GTX89955 ELMO1 antibody, C-term.

Dilution : 0.3 $\mu$ g/ml

Loading : 35 $\mu$ g protein in RIPA buffer

**GTX89955 WB Image**

WB analysis of Jurkat cell lysate using GTX89955 ELMO1 antibody, C-term.

Dilution : 0.3 $\mu$ g/ml

Loading : 35 $\mu$ g protein in RIPA buffer



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 03 Page 2 of 2